

BEST AVAILABLE COPY

Scan of the stained blotting membrane from Example 1



Lane	1	2	3	4	5	6	7	8
Final dilution	1 : 10	20	40	80	160	320	640	1280


Fig.1

BEST AVAILABLE COPY

Final dilution 1:10

- 1 Standard human plasma


Bands (%)	
1-5	40
6-10	30
11...	30



Final dilution 1:20

- 2 Standard human plasma


Bands (%)	
1-5	38
6-10	33
11...	29



Final dilution 1:40

- 3 Standard human plasma


Bands (%)	
1-5	36
6-10	31
11...	33



Final dilution 1:80

- 4 Standard human plasma


Bands (%)	
1-5	38
6-10	32
11...	30



Final dilution 1:160

- 5 Standard human plasma

Bands (%)	
1-5	41
6-10	35
11...	24



Final dilution 1:320

- 6 Standard human plasma

Bands (%)	
1-5	40
6-10	33
11...	27





Fig.2

Scan of the stained blotting membrane from Example 2



Lane	1	2	3	4	5	6	7	8
Final dilution (l/ml)		1:	0.2					
		2:	0.1					
		3:	0.05					
		4:	0.025					
		5:	0.0125					
		6:	0.006					
		7:	0.003					
		8:	0.0015					

Fig.3

BEST AVAILABLE COPY

Final dilution 0.2 IU/ml

1 Heamate HS

Bands (%)

1-5 47

6-10 35

11... 18

Final dilution 0.1 IU/ml

2 Heamate HS

Bands (%)

1-5 44

6-10 37

11... 18

Final dilution 0.05 IU/ml

3 Heamate HS

Bands (%)

1-5 42

6-10 39

11... 19

Final dilution 0.025 IU/ml

4 Heamate HS

Bands (%)

1-5 41

6-10 43

11... 16

Final dilution 0.0125 IU/ml

5 Heamate HS

Bands (%)

1-5 47

6-10 38

11... 16

Final dilution 0.006 IU/ml

6 Heamate HS

Bands (%)

1-5 81

6-10 3

11... 16









Fig.4

Scan of the Coomassie blue-stained gel from Example 3



BEST AVAILABLE COPY

- 5 Humate-P (150µl sample + 50µl buffer)
- 6 Humate-P (200µl sample + 50µl buffer)
- 7 Haemate-HS (150µl sample + 50µl buffer)
- 8 Haemate-HS (200µl sample + 50µl buffer)

Fig.5

Lane 5


Humate-P

Bands (%)

1-5 23

6-10 40

11... 38



Lane 6


Humate-P

Bands (%)

1-5 25

6-10 40

11... 36



Lane 7

Haemate-HS


Bands (%)

1-5 19

6-10 40

11... 41

(Evaluation without artefact)



Lane 8

Haemate-HS

Bands (%)

1-5 20

6-10 38

11... 43




Fig.6

Scan of the Coomassie blue-stained gel from Example 4




Fig.7

BEST AVAILABLE COPY

Scan of the immunostained blotting membrane from Example 5

Lanes 1 and 8: Fibrinogen reference sample

Lanes 2-7: various fibrinogen batches






Fig.8


BEST AVAILABLE COPY




1
Standard Human Plasma



2
Humate P



3
Humate-P



4
Humate-P

Fig.9